

Author index to volume 261

- Abel, R.J.R., N.J. Finizio and M. Greig, (3,6) GWhD(v)—existence results (1–3) 3–26
- Aldred, R.E.L., J. Širáň and M. Širáň, A note on the number of graceful labellings of paths (1–3) 27–30
- Anderson, I. and D.A. Preece, Power-sequence terraces for \mathbb{Z}_n where n is an odd prime power (1–3) 31–58
- Batten, L.M. and S. Sane, Permutations with a distinct difference property (1–3) 59–67
- Bennett, F.E., B. Du and H. Zhang, Existence of self-orthogonal diagonal Latin squares with a missing subsquare (1–3) 69–86
- Billington, E.J. and D.G. Hoffman, Decomposition of complete tripartite graphs into gregarious 4-cycles (1–3) 87–111
- Buratti, M. and A. Del Fra, Existence of cyclic k -cycle systems of the complete graph (1–3) 113–125
- Chang, Y. and Y. Miao, Constructions for optimal optical orthogonal codes (1–3) 127–139
- Colbourn, C.J. and A.C.H. Ling, Graph decompositions with application to wavelength add-drop multiplexing for minimizing SONET ADMs (1–3) 141–156
- Danziger, P. and E. Mendelsohn, Bicolour graphs of Steiner triple systems (1–3) 157–176
- Dejter, I.J., Equitable factorizations of Hamming shells (1–3) 177–187
- Del Fra, A., see M. Buratti (1–3) 113–125
- Deng, D., M. Greig and P.R.J. Östergård, More c -Bhaskar Rao designs with small block size (1–3) 189–208
- Deng, D., R. Rees and H. Shen, On the existence and application of incomplete nearly Kirkman triple systems with a hole of size 6 or 12 (1–3) 209–233
- de Resmini, M.J. and D. Jungnickel, Two infinite families of failed symmetric designs (1–3) 235–241
- Dinitz, J.H., E.R. Lamken and A.C.H. Ling, Complementary partial resolution squares for Steiner triple systems (1–3) 243–254
- Du, B., see F.E. Bennett (1–3) 69–86
- Finizio, N.J., see R.J.R. Abel (1–3) 3–26
- Forbes, A.D., M.J. Grannell and T.S. Griggs, On colourings of Steiner triple systems (1–3) 255–276
- Grannell, M.J., see A.D. Forbes (1–3) 255–276
- Gray, B.D., R. Mathon, T. Moran and A.P. Street, The spectrum of minimal defining sets of some Steiner systems (1–3) 277–284
- Gray, I.D., J. MacDougall, J.P. McSorley and W.D. Wallis, Vertex-magic labeling of trees and forests (1–3) 285–298
- Greig, M. and D.H. Rees, Existence of balanced incomplete block designs for many sets of treatments (1–3) 299–324
- Greig, M., see D. Deng (1–3) 189–208
- Greig, M., see R.J.R. Abel (1–3) 3–26
- Griggs, T.S., see A.D. Forbes (1–3) 255–276

- Hoffman, D.G., C.C. Lindner and C.A. Rodger, A partial $2k$ -cycle system of order n can be embedded in a $2k$ -cycle system of order $kn + c(k)$, $k \geq 3$, where $c(k)$ is a quadratic function of k (1-3) 325-336
- Hoffman, D.G., see E.J. Billington (1-3) 87-111
- Hurd, S.P. and D.G. Sarvate, An application of partition of indices to enclosings of triple systems (1-3) 337-346
- Ji, L. and L. Zhu, Constructions for Steiner quadruple systems with a spanning block design (1-3) 347-360
- Jungnickel, D. and L. Storme, Maximal partial spreads in $PG(3,4)$ and maximal sets of mutually orthogonal latin squares of order 16 (1-3) 361-371
- Jungnickel, D., see M.J. de Resmini (1-3) 235-241
- Kreher, D.L., A.C.H. Ling, R.S. Rees and C.W.H. Lam, A note on $\{4\}$ -GDDs of type 2^{10} (1-3) 373-376
- Lam, C.W.H., see D.L. Kreher (1-3) 373-376
- Lamken, E.R., see J.H. Dinitz (1-3) 243-254
- Lindner, C.C., see D.G. Hoffman (1-3) 325-336
- Ling, A.C.H., An application of splittable 4-frames to coloring of $K_{n,n}$ (1-3) 377-382
- Ling, A.C.H., see C.J. Colbourn (1-3) 141-156
- Ling, A.C.H., see D.L. Kreher (1-3) 373-376
- Ling, A.C.H., see J.H. Dinitz (1-3) 243-254
- Liu, Z. and H. Shen, Embeddings of almost resolvable triple systems (1-3) 383-398
- MacDougall, J., see I.D. Gray (1-3) 285-298
- Mathon, R., see B.D. Gray (1-3) 277-284
- McSorley, J.P., see I.D. Gray (1-3) 285-298
- Mendelsohn, E., see P. Danziger (1-3) 157-176
- Miao, Y., see Y. Chang (1-3) 127-139
- Milazzo, L., Z. Tuza and V. Voloshin, Strict colorings of Steiner triple and quadruple systems: a survey (1-3) 399-411
- Moran, T., see B.D. Gray (1-3) 277-284
- Östergård, P.R.J., see D. Deng (1-3) 189-208
- Preece, D.A., see I. Anderson (1-3) 31-58
- Quattrocchi, G., Embedding handcuffed designs in D -designs, where D is the triangle with attached edge (1-3) 413-434
- Rees, D.H., see M. Greig (1-3) 299-324
- Rees, R., see D. Deng (1-3) 209-233
- Rees, R.S., see D.L. Kreher (1-3) 373-376
- Rodger, C.A., see D.G. Hoffman (1-3) 325-336
- Sane, S., see L.M. Batten (1-3) 59-67
- Sarvate, D.G., see S.P. Hurd (1-3) 337-346
- Shalaby, N., The existence of near-Rosa and hooked near-Rosa sequences (1-3) 435-450
- Shen, H., see D. Deng (1-3) 209-233
- Shen, H., see Z. Liu (1-3) 383-398
- Širáň, J., see R.E.L. Aldred (1-3) 27-30
- Širáň, M., see R.E.L. Aldred (1-3) 27-30

- Storme, L., see D. Jungnickel (1-3) 361-371
Street, A.P., see B.D. Gray (1-3) 277-284

Tuza, Z., see L. Milazzo (1-3) 399-411

Voloshin, V., see L. Milazzo (1-3) 399-411

Wallis, W.D., see I.D. Gray (1-3) 285-298

Yin, J., A new proof of a Colbourn-Rosa problem (1-3) 451-458

Zhang, H., see F.E. Bennett (1-3) 69-86
Zhu, L., see L. Ji (1-3) 347-360

the 1980s, the number of people in the world who are illiterate has increased from 1.2 billion to 1.5 billion (UNESCO 1991).

There are a number of reasons why illiteracy has increased. First, the population of the world has increased by 1 billion in the last 20 years. Second, the number of people who are illiterate has increased in all regions of the world, although the rate of increase has been highest in sub-Saharan Africa and South Asia. Third, the number of people who are illiterate has increased in all age groups, although the rate of increase has been highest in the young adult population (UNESCO 1991).

There are a number of reasons why illiteracy has increased in all regions of the world. First, the number of people who are illiterate has increased in all regions of the world, although the rate of increase has been highest in sub-Saharan Africa and South Asia. Second, the number of people who are illiterate has increased in all age groups, although the rate of increase has been highest in the young adult population (UNESCO 1991).

There are a number of reasons why illiteracy has increased in all regions of the world. First, the number of people who are illiterate has increased in all regions of the world, although the rate of increase has been highest in sub-Saharan Africa and South Asia. Second, the number of people who are illiterate has increased in all age groups, although the rate of increase has been highest in the young adult population (UNESCO 1991).

There are a number of reasons why illiteracy has increased in all regions of the world. First, the number of people who are illiterate has increased in all regions of the world, although the rate of increase has been highest in sub-Saharan Africa and South Asia. Second, the number of people who are illiterate has increased in all age groups, although the rate of increase has been highest in the young adult population (UNESCO 1991).

There are a number of reasons why illiteracy has increased in all regions of the world. First, the number of people who are illiterate has increased in all regions of the world, although the rate of increase has been highest in sub-Saharan Africa and South Asia. Second, the number of people who are illiterate has increased in all age groups, although the rate of increase has been highest in the young adult population (UNESCO 1991).

There are a number of reasons why illiteracy has increased in all regions of the world. First, the number of people who are illiterate has increased in all regions of the world, although the rate of increase has been highest in sub-Saharan Africa and South Asia. Second, the number of people who are illiterate has increased in all age groups, although the rate of increase has been highest in the young adult population (UNESCO 1991).

